

Symmetry, Confinement and the phase diagram of QCD

Adriano Di Giacomo

Pisa University and INFN Sezione di Pisa

Abstract

A general discussion is presented of the possible symmetries responsible for confinement of color and of their evidence in lattice simulations. The consequences on the phase diagram of *QCD* are also analyzed.

Key words: Non perturbative *QCD*, Confinement, Deconfining transition, Duality.

PACS: 11.10Wx, 11.15Ha, 12. 38Mh, 64. 60Cn

1. Why symmetry?

No free quark has ever been observed in Nature: the abundance of quarks n_q compared to the abundance of protons n_p has an experimental upper bound $\frac{n_q}{n_p} \leq 10^{-27}$ to be compared to the value 10^{-12} expected in the Standard Cosmological Model in absence of confinement. The cross section for inclusive production of quarks in hadron collisions, σ_q is also 10^{-15} times smaller than the perturbative expectation. The natural explanation of these facts is that confinement is an absolute property, in the sense that n_q and σ_q are strictly zero due to some symmetry. As a consequence the deconfining transition is a change of symmetry, i.e. an order-disorder transition and can not be a cross-over. A similar situation exists in ordinary superconductivity: the resistivity in the superconducting phase has an exceedingly small experimental upper limit compared to the resistivity in the normal phase. The natural explanation is that the resistivity in the superconducting phase is strictly zero. A change of symmetry occurs at the transition from a Higgs broken $U(1)$ symmetry (superconductor) in which Cooper pairs condense in the vacuum, to a normal phase in which the $U(1)$ symmetry is exact.

2. What symmetry?

Color symmetry is exact : it can not distinguish confined from deconfined. Center symmetry only exists in absence of dynamical quarks. Chiral symmetry only exists at zero m_q . Moreover in some cases like QCD with $N_f = 2$ adjoint fermions chiral symmetry restoration occurs at a different temperature than deconfinement [1][2], indicating that the relevant degrees of freedom at the deconfining transition are not the chiral ones. The only way to get an extra symmetry is via duality [3][4], i.e. by looking at excitations with topologically non trivial boundary conditions. In $(2 + 1)dim$ the homotopy is Π_1 and the topologically non trivial excitations are vortices, in $(3 + 1)dim$ the homotopy is Π_2 and the excitations are monopoles [5][6]. For a generic gauge group G of rank r , r abelian field strength tensors ('t Hooft tensors) $F_{\mu\nu}^a$, ($a = 1, ..r$) can be defined[7] and in terms of them r magnetic currents $j_\nu^a \equiv \partial_\mu F_{\mu\nu}^{a*}$. Non zero value of the currents j_ν^a is a violation of Bianchi identities, due to the presence of magnetic charges. The currents j_ν^a are conserved due to the antisymmetry of the dual tensor $F_{\mu\nu}^{a*}$ and define the dual symmetry. If the corresponding $U(1)$ symmetries are Higgs broken magnetic charges condense in the vacuum and there is dual superconductivity (Confinement). If the symmetries are exact the vacuum is normal and chromoelectric charges deconfined. An operator μ can be constructed carrying non zero magnetic charge, and its *vev* $\langle\mu\rangle$ can be used as an order parameter for confinement [8][9][10], i.e. as a detector of monopole condensation.

3. The phase diagram.

A transition is a rapid change in physics at some value T_c of some parameter say the temperature T . A transition shows up as a peak in susceptibilities, which are the derivatives of observables with respect to T . For example a peak in the specific heat C_V is a rapid change in the heat content. A transition is called a crossover if no discontinuity develops at T_c as the volume V goes to infinity, it is named first order if some first derivative of the free energy diverges, e.g. if the free energy itself has a discontinuity at T_c and C_V diverges as $V \rightarrow \infty$.

Stating that a transition is a crossover is equivalent to verify that the free energy is analytic through T_c , and this cannot be done on the basis of any numerical calculations with a finite volume and a finite resolution. It can sometime be done with the help of some theory. A classical example is the chiral transition at small quark masses in $N_f = 2$ QCD [11]. Assuming that the relevant degrees of freedom at the chiral-deconfinement transition are the chiral ones, on the basis of renormalization group arguments one can say that either the chiral transition is second order in the universality class of $O(4)$, and then the transition is a cross-over at small non zero masses, or it is first order, and then it stays first order at small masses. In the first case a tricritical point is predicted at finite density, whose existence can be checked experimentally in heavy ion collisions[12]; no tricritical point exists in the second case. Finite size scaling analysis has been performed by many groups[13], but none finds evidence for second order $O(4)$.

If the correlation lengths are large compared to lattice spacing scale invariance holds and one expects for the volume dependence e.g. of the specific heat the following scaling law[14][15]

$$C_V - C_0 \approx L_s^{\frac{\alpha}{\nu}} \Phi_C(\tau L_s^{\frac{1}{\nu}}, m L_s^{y_h}) \quad (1)$$

Here L_s is the spatial size of the lattice, $\tau = (1 - \frac{T}{T_c})$ the reduced temperature and α , ν and y_h are critical indexes which are specific of the order and universality class of the transition. For second order $O(4)$ $\alpha = -.24$, $\frac{1}{\nu} = 1.34$, $y_h = 1.48$. For weak first order $\alpha = 1$, $\frac{1}{\nu} = 3$, $y_h = 3$. Eq.(1) can be tested on lattice data either by keeping the second variable of the function Φ_C fixed, by choosing m and L_s such that $m L_s^{y_h}$ has a fixed value, say K for a given assumption (y_h) on the universality class; or by keeping the first variable fixed and checking the dependence on the second one[14]. One has in the first case

$$(C_V - C_0)/L_s^{\frac{\alpha}{\nu}} \approx \Phi_C(\tau L_s^{\frac{1}{\nu}}, K) \quad (2)$$

in the second case, at large values of $m L_s^{y_h}$ [15], one has for second order $O(4)$

$$(C_V - C_0) \approx m^{.13} f_C(\tau L_s^{1.35}) \quad (3)$$

Instead for weak first order

$$(C_V - C_0) \approx L_s^3 f_C^0(\tau L_s^3) + \frac{1}{m} f_C^1(\tau L_s^3) \quad (4)$$

Data on lattices $L_t = 4$, $L_s = 16, 20, 24, 32$ do not agree with the scaling Eq.(2) with the choice $O(4)$, they do with the choice weak 1st order [14][15]. Also Eq(3) is not satisfied by $O(4)$. Eq.(4) instead is obeyed, but the first term, which is typical of first order looks to be negligible at present volumes, implying that the transition is too weak to observe a growth proportional to the volume at presently available volumes. Moreover, with $L_t = 4$ the lattice at the phase transition is rather coarse, and a check should be done with smaller lattice spacings. Evidence for the existence of the first term of Eq.(4) is needed to make a definite statement on first order. No definite evidence exists by now for a cross-over.

References

- [1] F. Karsch, M. Lutgemeier Nucl.Phys.**B550**, 449(1999)
- [2] G. Cossu , M.D'Elia, A. Di Giacomo , G. Lacagnina , C. Pica; Phys.Rev.**D77**:074506,2008.
- [3] H.A. Kramers, G.H. Wannier; Phys. Rev.**66**,252 (1941)
- [4] S. Gukov and E. Witten; Gauge theory, ramification, and the geometric Langlands program, [hep-th/0612073].
- [5] G. 't Hooft; Nucl. Phys. B **79** (1974) 276.
- [6] A. M. Polyakov; JETP Lett. **20** (1974) 194
- [7] A. Di Giacomo, L. Lepori, F. Pucci; JHEP 0810:096,2008.
- [8] A. Di Giacomo, G. Paffuti; Phys.Rev.D56:6816-6823,1997.
- [9] A. Di Giacomo, B. Lucini, L. Montesi, G. Paffuti ; Phys.Rev.D61:034503,2000.
- [10] A. Di Giacomo, B. Lucini, L. Montesi, G. Paffuti ; Phys.Rev.D61:034504,2000.
- [11] R.D. Pisarski, F. Wilczek ; Phys. Rev.**D29**, 338 (1984)
- [12] M.A. Stephanov, K. Rajagopal, E.V. Shuryak; Phys.Rev.Lett. 81, 4816 (1998)
- [13] See e.g. Owe Philipsen , Status of Lattice Studies of the QCD Phase Diagram. International Symposium Fundamental Problems in Hot and / or Dense QCD, Kyoto, Japan -2008. arXiv:0808.0672 [hep-ph]

- [14] M. D'Elia, A. Di Giacomo, C. Pica, Phys.Rev.D72:114510,2005.
- [15] G.Cossu, M. D'Elia, A. Di Giacomo, C. Pica, arXiv:0706.4470 [hep-lat]